



## Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

*Demonstration.*—Put  $OP_m$  to represent the  $m^{th}$  of the  $n$  roots of  $OP$ , then if the proposition is true  $OP_m^n = OP$ .

By prop. 1  $OP_m^n = OR$  revolved through  $n(\frac{1}{n}\theta + \frac{1}{n}(m-1)C)$ ,  
or  $OP_m^n = OR$  “ “ “  $\theta$ ,

since complete circumferences may be rejected.

But  $OP = OR$  revolved through  $\theta$ , therefore  $OP_m^n = OP$ .

PROPOSITION 4.— $OR^{\frac{1}{n}} = OR$  revolved through  $\frac{1}{n}C$ .

Hence when  $OP = OR^{\frac{1}{n}}$ ,  $OP$  is the  $n^{th}$  root of  $+1$ , or  $OR$ .

*Demonstration.*—When  $\theta = RP$  becomes  $C$ ,  $OP$  becomes  $OR$ , hence by substitution  $OP^{\frac{1}{n}} = OR$  revolved through  $\frac{1}{n}C$  (prop. 2),

becomes  $OR^{\frac{1}{n}} = OR$  “ “ “  $\frac{1}{n}C$ .

This formula gives but one of the  $n$  roots of  $+1$ , or  $OR$ , which root I call the first of the  $n$  roots of  $+1$ .

By prop. 3 the  $m^{th}$  of the  $n-1$  remaining roots of  $+1$ , is the  $m^{th}$  power of the  $n^{th}$  root of  $+1$ .

It is plain that the  $n$  roots of  $+1$  are *numerically* equal to each other but *geometrically* or *positionally* different.

The reader may test the above propositions by solving the binomial equations  $x^2 = 1$ ,  $x^3 = 1$ ,  $x^4 = 1$ ,  $x^5 = 1$ ,  $x^6 = 1$ ,  $x^8 = 1$ ,  $x^{10} = 1$ ,  $x^{12} = 1$ ,  $x^{20} = 1$ , etc., and constructing the roots in accordance with the convention that  $\sqrt{-1}$  denotes *perpendicularity*. See my first article in the January ANALYST The passage from  $OR = 1$  to  $OR = a$  is simple. The next article will deal with imaginary exponents.

## ~~~~~

### SOLUTION OF AN INDETERMINATE PROBLEM.

BY GEO. R. PERKINS, L. L. D., UTICA, N. Y.

*No. 5.—Find three right triangles having equal perimeters, and whose areas shall be in arithmetical progression.*

Since these triangles have the same perimeters, and their areas are in arithmetical progression, it is evident that the radii of their inscribed circles must be in arithmetical progression. We will therefore assume as follows:

$$\text{First triangle. } \begin{cases} r + s & = \text{radius of inscribed circle.} \\ x + (r + s) & = \text{the base.} \\ y + (r + s) & = \text{the perpendicular.} \end{cases}$$

$$\text{Second triangle. } \begin{cases} r & = \text{radius of inscribed circle.} \\ x' + r & = \text{the base.} \\ y' + r & = \text{the perpendicular.} \end{cases}$$

$$\text{Third triangle. } \begin{cases} r - s & = \text{radius of inscribed circle.} \\ x'' + (r - s) & = \text{the base.} \\ y'' + (r - s) & = \text{the perpendicular.} \end{cases}$$

Denoting the common perimeter of the three triangles by  $p$ , we have for their hypotenuses as follows:

$$x + y = \frac{1}{2}p - (r + s) \dots\dots\dots(1),$$

$$x' + y' = \frac{1}{2}p - r \dots\dots\dots(2),$$

$$x'' + y'' = \frac{1}{2}p - (r - s) \dots\dots\dots(3).$$

These conditions give  $(x + y) + (x'' + y'') = 2(x' + y')$ , which proves the hypotenuses to be in arithmetical progression.

Since the triangles are right, we have

$$[x + (r + s)]^2 + [y + (r + s)]^2 = (x + y)^2 \dots\dots\dots(4),$$

$$[x' + r]^2 + [y' + r]^2 = (x' + y')^2 \dots\dots\dots(5),$$

$$[x'' + (r - s)]^2 + [y'' + (r - s)]^2 = (x'' + y'')^2 \dots\dots\dots(6).$$

These, when simplified, become

$$xy = (r + s)(x + y) + (r + s)^2 \dots\dots\dots(7),$$

$$x'y' = r(x' + y') + r^2 \dots\dots\dots(8),$$

$$x''y'' = (r - s)(x'' + y'') + (r - s)^2 \dots\dots\dots(9).$$

Using the values of (1), (2) and (3), we have

$$xy = \frac{1}{2}p(r + s) \dots\dots\dots(10),$$

$$x'y' = \frac{1}{2}pr \dots\dots\dots(11),$$

$$x''y'' = \frac{1}{2}p(r - s) \dots\dots\dots(12).$$

If from the squares of (1), (2) and (3) we subtract four times these products, and extract the square roots, we find

$$x - y = \sqrt{\frac{1}{4}p^2 - 3p(r + s) + (r + s)^2} \dots\dots\dots(13),$$

$$x' - y' = \sqrt{\frac{1}{4}p^2 - 3pr + r^2} \dots\dots\dots(14),$$

$$x'' - y'' = \sqrt{\frac{1}{4}p^2 - 3p(r - s) + (r - s)^2} \dots\dots\dots(15).$$

Hence, we must have

$$\frac{1}{4}p^2 - 3p(r + s) + (r + s)^2 = \square \dots\dots\dots(16),$$

$$\frac{1}{4}p^2 - 3pr + r^2 = \square \dots\dots\dots(17),$$

$$\frac{1}{4}p^2 - 3p(r - s) + (r - s)^2 = \square \dots\dots\dots(18).$$

Condition (16) will be satisfied if we take  $p = 12(r + s)$ , and (17) and (18) will become

$$r^2 + 36rs + 36s^2 = \square \dots\dots\dots(19),$$

$$r^2 + 70rs + 73s^2 = \square \dots\dots\dots(20).$$

Conditions (19) and (20) will be satisfied, by taking  $r = 9s$ , and consequently  $p = 120s$ .

Taking  $s = 1$ , we immediately obtain

$$x + y = 50; \quad x' + y' = 51; \quad x'' + y'' = 52;$$

$$x - y = 10; \quad x' - y' = 21; \quad x'' - y'' = 28.$$

Consequently,  $x = 30; \quad y = 20;$

$$x' = 36; \quad y' = 15;$$

$$x'' = 40; \quad y'' = 12.$$

$$\text{First triangle.} \quad \left\{ \begin{array}{ll} x + r + s = 40 = \text{base.} \\ y + r + s = 30 = \text{perpendicular.} \\ x + y = 50 = \text{hypotenuse.} \\ \hline 120 = \text{perimeter.} \\ 600 = \text{area.} \end{array} \right.$$

$$\text{Second triangle.} \quad \left\{ \begin{array}{ll} x' + r = 45 = \text{base.} \\ y' + r = 24 = \text{perpendicular.} \\ x' + y' = 51 = \text{hypotenuse.} \\ \hline 120 = \text{perimeter.} \\ 540 = \text{area.} \end{array} \right.$$

$$\text{Third triangle.} \quad \left\{ \begin{array}{ll} x'' + r - s = 48 = \text{base.} \\ y'' + r - s = 20 = \text{perpendicular.} \\ x'' + y'' = 52 = \text{hypotenuse.} \\ \hline 120 = \text{perimeter.} \\ 480 = \text{area.} \end{array} \right.$$

I presume these results are the least values which will satisfy our Problem.

REMARK. —This Problem is the same as Problem 103, on page 465

of *John D. Williams' Elementary Treatise on Algebra*, Boston, 1840. The solution there given is satisfactory, but extremely lengthy, and very complex. The results obtained are as follows:

The sides of the first triangle      18601944; 13951458; 23252430.  
 “   “   “   “ second triangle    13999464; 18559223; 23247145.  
 “   “   “   “ third triangle      18515584; 14048388; 23241860.

Common perimeter = 55805832.

Radii of the inscribed circles are 4650486; 4655771; 4661056.

### ~~~~~

*SOLUTION OF A PROBLEM.*

~~~~~

BY PROF. DAVID TROWBRIDGE, WATERBURGH, N. Y.

Required the Sum of the Products of  $n$  Quantities taken  $m$  and  $m$  together, no two products having the same Factors.

Let the  $n$  quantities be  $r_1, r_2, r_3, \dots, r_n$ . Let  $S_m$  be the required sum. Also let

$$S_1 = R_1 = r_1 + r_2 + \dots + r_n, \quad R_2 = r_1^2 + r_2^2 + \dots + r_n^2, \dots$$

$$R_m = r_1^m + r_2^m + \dots + r_n^m \quad (1).$$

Now

$$r_1(r_2 + r_3 + \dots + r_n) + r_2(r_1 + r_3 + \dots + r_n) + \dots$$

$$+ r_n(r_1 + r_2 + \dots + r_{n-1}) = r_1(R_1 - r_1) + r_2(R - r_2) + \dots$$

$$+ r_n(R_1 - r_n) = R_1^2 - R_2 = R_1 S_1 - R_2.$$

But we have evidently taken each product twice, so that we shall consequently have

$$2S_2 = R_1 S_1 - R_2 \dots \dots \dots (2).$$

$$r_1(r_2 r_3 + r_2 r_4 + \dots) + r_2(r_1 r_3 + r_1 r_4 + \dots) + \dots$$

$$+ r_n(r_1 r_2 + r_1 r_3 + \dots) = r_1[S_2 - r_1(R_1 - r_1)]$$

$$+ r_2[S_2 - r_2(R_1 - r_2)] + \dots + r_n[S_2 - r_n(R_1 - r_n)]$$

$$= R_1 S_2 - R_1 R_2 + R_3 = R_1 S_2 - R_2 S_1 + R_3.$$

But we have now taken each product three times. We hence have

$$3S_3 = R_1 S_2 - R_2 S_1 + R_3 \dots \dots \dots (3).$$